



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

of lactation. But few cows change one per cent from richest milk of last month before drying up to thinnest milk after calving.

The following is the record of six cows at the Experiment Station Farm that calved in the spring and were fed at the barn heavily with grain, hay, and ensilage, before and during pasturage, and also after their return to the barn until they dried up.

	April.	May.	June.	July.	August.	September.	October.	November.
Average monthly yield per cow, pounds.....	792	867	918	814	723	711	531	340
Ratio of different months, if June is 100.....	84	91	100	86	76	75	56	36
Average per cent of fat in milk.....	4.07	4.38	4.38	4.38	4.37	4.52	4.70	4.83
Ratio of different months, if June is 100.....	93	100	100	98	100	103	107	110
Average monthly yield of butter-fat per cow, lbs...	32.2	38.0	41.5	35.8	31.6	32.1	25.0	16.4
Ratio of different months, if June is 100.....	78	91	100	84	76	77	60	40

The influence of full feeding is seen most strongly during the months of April and May, which yield, with grain, one-third more milk and butter-fat than without. An influence after June is seen, but not so pronounced. Those having grain shrink in milk-flow only nine-tenths as fast as those not having grain, and have the advantage of only one-twenty-fifth in the shrinkage of butter-fat.

Of course, this is not a strict comparison of the effects of feeding grain on the total yield or of the financial side of the question, but merely of the effect the grain has of increasing the flow of the milk at once when the cow calves and of maintaining the milk-flow for a longer period in the latter part of lactation.

#### LETTERS TO THE EDITOR.

\*\*\* Correspondents are requested to be as brief as possible. The writer's name is in all cases required as proof of good faith.

On request in advance, one hundred copies of the number containing his communication will be furnished free to any correspondent.

The editor will be glad to publish any queries consonant with the character of the journal.

#### An Unusual Aurora.

ON Saturday evening, July 15, there occurred an aurora which was unlike any the writer has ever seen, and a brief description of it may contribute something to the aggregate knowledge of those interesting phenomena.

The peculiar feature of this aurora was the movement of a series or succession of whitish flecks across the sky from east to west, resembling somewhat the waves of a body of water.

About 9.30, central time, my attention was first attracted to it. Flecks of white light were forming in the east at an altitude of about 45°, passing in regular succession westward, about 20° north of the zenith, and apparently accumulating in one larger band in the northwest, reaching at times from near the horizon to perhaps 80°. The white flecks or streaks were about 10° in length, strictly parallel north and south, and quite uniform in distance apart. They grew brighter and more distinct as they approached and passed the meridian. Their motion was very regular and quite rapid,—comparable to the swiftest apparent motion of light clouds. If they were as high as the electric theory would suggest, the velocity must have been enormous.

At times similar short bands, like strokes with a paint-brush, were stationary in the north, at about 45° altitude, for several minutes at a time.

A few minutes later a number, perhaps ten or twelve, white

bands appeared north of the zenith, all converging towards a point some 10° south of the zenith, but vanishing before reaching the zenith. They remained only a few minutes. About 10 o'clock the moving flecks had disappeared, and one long, straight band extended from the northwest horizon, 50° or 60°, toward a point about 45° south of the zenith. Two or three other short flecks appeared parallel with the main band. About the same time the usual diffused glow appeared in the north horizon and continued till after 11 o'clock, but was not observable while the moving bands were seen. Many more gorgeous auroras have been seen in our latitude, but the rapidly-moving bands gave this one a new interest.

W. H. HOWARD.

Adrian, Mich.

#### Light-Shunners and Light-Seekers.

It is well known that in the main divisions of the animal world we find groups which normally withdraw from daylight and which form a very large minority of existing species. Some of these lovers of darkness dwell in caverns, in underground burrows or in the seas at depths where the light penetrates feebly or not at all.

We might, perhaps, expect that such creatures would feel annoyed, more or less, by artificial light and would withdraw from what to them must be an exceptional phenomenon. This, however, would be a mistake. The only nocturnal animals which seem to shun fire and light are the carnivorous mammals especially the cats. It has long been customary for travellers in Africa to keep lions, leopards, etc., aloof from an encampment by means of bonfires. As a rule the sleepers are safe as long as the fires are fed up.

The lemurs and loris are even more nocturnal than the cats, since they do not travel or prey by day. Whether they are repelled or attracted by a light is not sufficiently decided.

The bats are not purely nocturnal. They are sometimes seen hawking for insects in full daylight. But a light attracts them. Entomologists—I may mention Major Elwes, P. E. S.—who have hung out lamps in order to entice moths, have often found that bats come to the lights and secure a large share of the specimens.

Among birds there are few truly nocturnal species. The owl and the night-jar (absurdly called the goat-sucker) are the most common night fliers. The owls are attracted by a light, a fact which has given rise to a foolish superstition. They will often dash against the window of a room which is lighted up by night. If, as often happens to be the case, this is a sick-chamber, nurses of the old school pronounce such a visit a fatal omen. Some would-be wise men have gravely asserted that the owl scents the approach of dissolution and comes in the hope of feasting upon the corpse. Now, in fact, the owl feeds by preference on prey which it has just killed, and in captivity it rejects any food which is in the slightest degree tainted.

In Australia the emur, though not truly nocturnal, may be seen rapidly scudding over the plains by moonlight.

Many birds which are perfectly diurnal, in their ordinary habits, fly by night when migrating, and are then attracted by a light. Numbers of various species dash themselves against the windows of lighthouses and are killed by the shock. This is much to be regretted, since the majority of migratory birds feed on insects, and had they survived they would during the coming season have been hard at work ridding our crops of vermin.

The habits of reptiles vary greatly. The few European snakes, e.g., the viper, the asp, the Austrian adder, the grass snake and *Coronella lavis*, are rarely met with save in the brightest hours of the day. But of the African, Indian and Australian species it may be said:

"The snake that loves the twilight has come out, beautiful, still and deadly"—though they also bask in the sun. Nor are they scared away by lights or fire. One species, indeed, if it espies a fire in the forest, seeks to dash or drag the sticks away. Toads, newts and salamanders live very contentedly in the dark, but seem to regard a light with indifference.

The majority of fishes and other dwellers in the waters are decidedly attracted by lights.